Catalog Number: 230-30238



## Recombinant Monkeypox Virus EEV Envelope Glycoprotein, A35R

#### Source

Species Monkeypox Virus (MPXV)

Accession Number Q8V4U4

Gene Symbol A35R

Expressed Region Gln59-Thr181

**Synonyms** EEV Envelope Glycoprotein, A35R

#### **Preparation**

Expression System Human embryonic kidney 293 (HEK293) cells

Tag C-terminal His-tag

**Purification** His-tag affinity purification by immobilized metal ion affinity chromatography (IMAC)

Purity >95%

Purity Determined By SDS-PAGE under reducing conditions and visualized by Coomassie blue staining

Recombinant protein product has a calculated molecular mass of ?14 kDa. Due to the abundant glycosylation, it migrates as one major ?14 kDa protein band (non-glycosylated form) and slight increased smear bands (glycosylated form) in SDS-PAGE under DTT and beta-mercaptoethanol

reducing conditions.

### **Protein Specifications**

**Molecular Weight** 

Format Liquid

Formulation Filtered solution in PBS with 1% mannitol and 5% trehalose

**Concentration** Determined by BCA protein assay

Endotoxin Level 0.5 EU per µg of the protein as determined by the LAL method

**Recommended Applications**Functional Assay, Protein-protein Interaction, Post-translational Modifications, ELISA, EIA, Western Blotting, Dot Blotting, Immunoprecipitation, Protein Array, etc.

SDS-PAGE Image SDS-PAGEund or type unknown

Figure 1. Deglycosylation analysis of purified recombinant proteins. The same amount of purified proteins were untreated (Lane 2) or treated with protein deglycosylation enzymes under native (Lane 3) or reducing (Lane 4) conditions. Deglycosylation treatment resulted in a mobility shift of the protein to produce one reduced band at the expected size, thus indicating that the untreated

recombinant protein (Lane 2) was glycosylated.

Lane 1: protein standard ladder (kDa).

Lane 2: untreated protein.

Lane 3: treated protein with deglycosylation enzymes under native conditions. Lane 4: treated protein with deglycosylation enzymes under denature conditions.

## **Shipping**



# Storage/Stability

Upon arrival, the protein may be stored for 2 weeks at 4 °C. For long term storage, it is recommended to store at -20 °C or -80 °C in appropriate aliquots. Avoid repeated freeze-thaw cycles.

This product is furnished for LABORATORY RESEARCH USE ONLY.

Not for diagnostic or therapeutic use.

